

## Product datasheet for **PP1223B2**

### Galectin 1 (LGALS1) Rabbit Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	<b>Direct ELISA:</b> To detect Human Galectin-1 by direct ELISA (using 100 µl/well antibody solution), this antibody can be used at a concentration of ~1.0 µg/ml. This Biotin conjugated antibody allows the detection of at least 0.2-0.4 ng/well of recombinant Human Galectin-1. <b>Sandwich ELISA:</b> To detect Human Galectin-1 by Sandwich ELISA (using 100 µl/well antibody solution), this antibody can be used at a concentration of 0.25-1.0 µg/ml. This Biotin conjugated antibody, in conjunction with Purified anti-Human Galectin-1 antibody (PP1223P) as Capture antibody, allows the detection of at least 0.2-0.4 ng/well of recombinant Human Galectin-1. <b>Western blot:</b> To detect Human Galectin-1 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hGalectin-1 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	E.coli derived recombinant Human Galectin-1.
Specificity:	Reacts with Galectin-1.
Formulation:	PBS, pH 7.2 without preservatives. Label: Biotin State: Lyophilized purified Ig fraction. Label: conjugated
Reconstitution Method:	Restore in sterile PBS solution containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.
Purification:	Affinity Chromatography.
Conjugation:	Biotin
Storage:	Store the antibody prior to reconstitution at -20°C. Following reconstitution the antibody can be stored at 2-8°C for one month or at -20°C for longer. Avoid repeated freezing and thawing.



[View online »](#)

<b>Stability:</b>	Shelf life: one year from despatch.
<b>Gene Name:</b>	galectin 1
<b>Database Link:</b>	<a href="#">Entrez Gene 3956 Human P09382</a>
<b>Background:</b>	The galectins are a family of beta-galactoside-binding proteins implicated in modulating cell-cell and cell-matrix interactions. Galectin 1 may act as an autocrine negative growth factor that regulates cell proliferation. It also may regulate cell apoptosis and cell differentiation. It binds beta-galactoside, CD45, CD3 and CD4. Galectin 1 inhibits CD45 protein phosphatase activity and therefore the dephosphorylation of Lyn kinase. It is found in cardiac, smooth, and skeletal muscle, neurons, thymus, kidney, placenta and hematopoietic cells.
<b>Synonyms:</b>	LGALS1, Galectin1, Lectin galactoside-binding soluble 1, Beta-galactoside-binding lectin L-14-I, Lactose-binding lectin 1, S-Lac lectin 1, Galaptin, 14 kDa lectin, HPL, HBL, MAPK-activating protein PM12