

Product datasheet for **R1064**

IacZ Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, IHC, WB
Recommended Dilution:	Suitable for Immunoblotting (Western or Dot blot), ELISA, Immunoprecipitation and most immunological methods requiring high titer and specificity. Recommended Dilutions: ELISA: 1/65,000. Western blot: 1/500-1/2,000. IHC: 1/500.
Reactivity:	Escherichia coli
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Beta-galactosidase from E. coli
Specificity:	Assay by immunoelectrophoresis resulted in a single precipitin arc against purified and partially purified Beta Galactosidase [E. coli]. Cross reactivity against Beta Galactosidase from other sources may occur but have not been specifically determined.
Formulation:	0.02 M Potassiumphosphate, 0.15 M Sodium chloride, pH 7.2 with 0.01% Sodium azide as preservative State: Serum State: Lyophilized serum
Reconstitution Method:	Restore with 2.0 ml of deionized water (or equivalent).
Concentration:	lot specific
Purification:	Purified from monospecific antiserum by Delipidation and Defribination.
Conjugation:	Unconjugated



[View online »](#)

Storage:	<p>Store vial at 2-8°C prior to restoration. Centrifuge product if not completely clear after standing at room temperature. This product is stable for one month at 2-8°C as an undiluted liquid.</p> <p>For extended storage aliquot contents and freeze at -20°C or below. Avoid cycles of freezing and thawing.</p> <p>Dilute only prior to immediate use.</p>
Stability:	Shelf life: one year from despatch.
Database Link:	P00722
Background:	<p>Beta-galactosidase is coded by a gene (lac z) in the lac operon of Escherichia coli. It is a metalloenzyme that splits lactose into glucose and galactose. It hydrolyzes terminal, non-reducing beta-D-galactose residues in beta-D-galactosides.</p> <p>Activation by cations seems to be substrate dependent. K⁺, Na⁺, NH₄⁺, Rb⁺, Cs⁺ and Mn⁺⁺ all activate enzyme activity based upon the substrate used.</p>
Synonyms:	Beta-Gal tag, lacZ tag, b0344, JW0335, Beta-Gal Fusion Protein, Lactase

Product images:




Figure 1. Western blotting using Fluorescein conjugated anti-Beta-galactosidase antibody shows a band at ~117 kDa (lanes 1 - 3) corresponding to 60 ng, 30 ng and 15 ng, respectively of b-Gal present in partially purified preparations (arrowhead). Lane 4 shows no cross reactivity with proteins present in a non-specific control E. coli lysate. Proteins were resolved on a 4-20% Tris-glycine gel by SDS-PAGE and transferred to Nitrocellulose and blocking using blocking buffer for fluorescent Western blotting. The membrane was probed with Fluorescein conjugated anti-Beta-galactosidase (. F) diluted to 1/10,000. Reaction occurred for 2 hours at room temperature. Molecular weight estimation was made by comparison to a prestained MW marker in lane M. Fluorescence image was captured using the VersaDoc® Imaging System. Other detection systems will yield similar results.