

## Product datasheet for R1074

### Choline oxidase Goat Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	Suitable for Immunoblotting (Western or Dot blot), ELISA, Immunoprecipitation and most immunological methods requiring high titer and specificity. <u>Recommended Dilutions:</u> This product has been assayed against 1.0 ug of Choline Oxidase [Alcaligenes species] in a standard sandwich ELISA using Peroxidase conjugated Affinity Purified anti-Goat IgG [H&L] (Rabbit) and ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:4,000 to 1:32,000 of the reconstitution concentration is suggested for this product.
Reactivity:	Alcaligenes
Host:	Goat
Clonality:	Polyclonal
Immunogen:	Choline Oxidase [Alcaligenes species].
Specificity:	Assay by immunoelectrophoresis resulted in a single precipitin arc against purified and partially purified Choline Oxidase [Alcaligenes species]. Cross reactivity against Choline Oxidase from other sources is unknown.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 0.01% sodium azide as preservative. State: Serum State: Lyophilized Serum
Reconstitution Method:	Restore with 2.0 ml of deionized water (or equivalent).
Concentration:	lot specific
Purification:	Prepared from monospecific antiserum by a delipidation and defibrination.
Conjugation:	Unconjugated



[View online »](#)

<b>Storage:</b>	Store vial at 2-8°C prior to restoration. Centrifuge product if not completely clear after standing at room temperature. For extended storage aliquot contents and freeze at -20°C or below. This product is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid cycles of freezing and thawing.
<b>Stability:</b>	Shelf life: One year from despatch.
<b>Database Link:</b>	<a href="#">P16101</a>
<b>Background:</b>	Choline Oxidase catalyzes the oxidation of choline and betaine aldehyde to betaine with concomitant consumption of oxygen and production of hydrogen peroxide.