

Product datasheet for R1075HRPS

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cholesterol Oxidase Goat Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, IP, WB

Recommended Dilution: Western blot: 1/500-1/5,000.

Immunoprecipitation: 1/100.

ELISA: 1/5,000-1/20,000.

This product has been assayed against 1.0 μ g of Cholesterol oxidase [microorganism] in a standard capture ELISA using ABTS as a substrate for 30 minutes at room temperature. A working dilution of 1/1,000 to 1/5,000 of the reconstitution concentration is suggested.

Reactivity: Bacteria **Host:** Goat

Clonality: Polyclonal

Immunogen: Cholesterol oxidase from Streptomyces

Specificity: This product detects Cholesterol oxidase (microorganism). Cross reactivity against Cholesterol

oxidase from other sources is unknown.

Immunoelectrophoresis give a single precipitin arc against anti-peroxidase, anti-goat serum

as well as purified and partially purified Cholesterol oxidase [microorganism].

Formulation: 0.02 M Potassium phosphate, 0.15 M Sodium chloride, pH 7.2

Label: HRP State: Purified

State: Lyophilized purified Ig fraction

Stabilizer: 10 mg/ml BSA (immunoglobulin and protease free)

Preservative: 0.01% (w/v) Gentamicin sulfate (Do NOT add Sodium azide!)

Label: Horseradish peroxidase

Reconstitution Method: Restore with 0.1 ml of deionized water (or equivalent).

Concentration: lot specific

Purification: Delipidation, salt fractionation and ion exchange chromatography followed by extensive

dialysis against the buffer

Conjugation: HRP





Cholesterol Oxidase Goat Polyclonal Antibody - R1075HRPS

Storage: Store lyophilized at 2-8°C for 6 months or at -20°C long term.

After reconstitution store the antibody undiluted at 2-8°C for one month or (in aliquots) at -

20°C long term.

Avoid repeated freezing and thawing.

Stability: Shelf life: one year from despatch.

Background: Cholesterol oxidases exist as both type I and type II oxidases and are implicated in bacterial

pathogenesis. In addition, they are important as clinical reagents, potential larvicides, and

tools in cell biology.

Synonyms: Cholesterol oxidase, CHOD, EC 1.1.3.6