

Product datasheet for R1091HRP

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GFP (Ads. to Hu, Ms, Rt Serum Proteins) Goat Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, IHC, WB

Recommended Dilution: Polyclonal anti-GFP antibody is designed to detect GFP and its variants. This antibody can be

used to detect GFP by ELISA (sandwich or capture) for the direct binding of antigen and recognizes wild type, recombinant and enhanced forms of GFP. Biotin conjugated polyclonal anti-GFP antibody used in a sandwich ELISA is well suited to titrate GFP in solution using this antibody in combination with a monoclonal anti-GFP antibody (Cat#R1461P) using either

form of the antibody as the capture or detection antibodies.

However, use the monoclonal form only for the detection of wild type or recombinant GFP as this form does not sufficiently detect 'enhanced' GFP. The detection antibody is typically conjugated to biotin and subsequently reacted with streptavidin conjugated HRP. Fluorochrome conjugated polyclonal anti-GFP antibody can be used to detect GFP by

immunofluorescence microscopy in prokaryotic (E.coli) and eukaryotic (CHO cells) expression systems and can detect GFP containing inserts. Significant amplification of signal is achieved using fluorochrome conjugated polyclonal anti-GFP antibody relative to the fluorescence of

GFP alone.

For Immunoblotting use either Alkaline Phosphatase or Peroxidase conjugated polyclonal

anti-GFP antibody to detect GFP or GFP containing proteins on western blots.

Recommended Dilutions: ELISA: 1/10,000-1/50,000. Western blot: 1/2,000-1/5,000.

Immunohistochemistry: 1/500-1/2,000.

Host: Goat

Clonality: Polyclonal

Immunogen: GST- Green Fluorescent Protein (GFP) fusion protein corresponding to the full length amino

acid sequence (246aa) derived from the jellyfish Aeguorea victoria

Specificity: Assay by Immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum,

anti-Peroxidase and purified and partially purified Green Fluorescent Protein (Aequorea

victoria).

No reaction was observed against Human, Mouse and Rat Serum Proteins.





GFP (Ads. to Hu, Ms, Rt Serum Proteins) Goat Polyclonal Antibody - R1091HRP

Formulation: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, containing 10 mg/ml BSA (IgG

and Protease free) as a stabilizer and 0.01% (w/v) Gentamicin Sulfate as a preservative.

Label: HRP

State: Lyophilized purified Ig fraction Label: Horseradish Peroxidase

Reconstitution Method: Restore with 1.0 ml of deionized water or equivalent.

Concentration: lot specific

Purification: Immunoaffinity Chromatography using Green Fluorescent Protein (Aequorea victoria) coupled

to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities.

Conjugation: HRP

Storage: Store vial at 2-8°C prior to restoration.

Restore with deionized water or equivalent; centrifuge product if not completely clear after

standing at room temperature.

This product is stable for one month at 2-8°C as an undiluted liquid. For extended storage aliquot contents and freeze at -20°C or below.

Avoid repeated freezing and thawing. Dilute only prior to immediate use.

Stability: Shelf life: one year from despatch.

Database Link: P42212

Background: Green fluorescence protein (GFP) is a 27 kDa protein derived from the jellyfish Aequorea

victoria, which emits green light (emission peak at a wavelenth of 509 nm) when excited by blue light (excitation peak at a wavelenth of 395 nm). Green Fluorescent Protein (GFP) has become an invaluable tool in cell biology research, since its intrinsic fluorescence can be visualized in living cells. GFP fluorescence is stable under fixation conditions and suitable for a variety of applications. GFP has been widely used as a reporter for gene expression, enabling researchers to visualize and localize GFP-tagged proteins within living cells without

between proteins through fluorescence energy transfer (FRET) protocols. GFP technnology has considerably contributed to a greater understanding of cellular physiology.

YFP differs from GFP due to a mutation at T203Y; antibodies raised against full-length GFP

the need for chemical staining. Other applications of GFP include assessment of protein protein interactions through the yeast two hybrid system and measurement of distance

should also detect YFP and other variants.

Synonyms: Green fluorescent protein, GFP-Tag

Note: Do **Not** Add Sodium Azide.