

## Product datasheet for R1091T

## Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

CN: techsupport@origene.cn

**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

## **GFP Goat Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** ELISA, IF, WB

Recommended Dilution: ELISA (Sandwich or Capture; ELISA for direct binding recognizes wild type, recombinant and

enhanced forms of GFP; for sandwich ELISA titrate GFP in solution using either form of the antibody as the capture or detection antibodies, detection antibody is typically conjugated to

to biotin and complexed with streptavidin-HRP).

Western blot (Peroxidase conjugated anti-GFP antibody shows a 42 kDa band).

Immunoflourescence.

**Host:** Goat

Clonality: Polyclonal

**Immunogen:** GST-Green Fluorescent Protein (GFP) fusion protein corresponding to the full length amino

acid sequence (246 aa) derived from the jellyfish Aequorea victoria

**Specificity:** This antibody detects Fluorescent Protein (GFP).

Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, anti-Peroxidase and purified and partially purified Green Fluorescent Protein (Aequorea victoria) Serum. No reaction was observed against Human, Mouse and Rat Serum Proteins.

**Formulation:** 0.125 M Sodium Borate, 0.075 M Sodium Chloride, 0.005 M EDTA, pH 8.0 as a buffer, 10

mg/ml Bovine Serum Albumin (BSA) IgG and (protease free) as stabilizer, 0.01 % (w/v) Sodium

azide as preservative

Label: TRITC

State: Lyophilized Ig fraction

Label: (Molecular weight 444 daltons) Absorption emission: 550 nm / 570 nm

Molar radio: 2.9 moles TRITC per mole of Goat IgG

**Reconstitution Method:** Restore with 1.0 ml of deionized water (or equivalent).

**Concentration:** lot specific

Purification: Immunoaffinity chromatography using Green Fluorescent Protein (Aequorea victoria) coupled

to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities

Conjugation: TRITC



## **GFP Goat Polyclonal Antibody - R1091T**

**Storage:** Store vial at 2 - 8 °C prior to reconstitution.

Following reconstitution the product is stable for one month at 2 - 8 °C as an undiluted liquid.

Dilute only prior to immediate use.

For extended storage add glycerol to 50 % and then aliquot contents and freeze at -20 °C or

below. Avoid repeated freezing and thawing.

Centrifuge product if not completely clear after standing at room temperature.

**Stability:** Shelf life: One year from despatch.

Database Link: P42212

**Background:** Green fluorescence protein (GFP) is a 27 kDa protein derived from the jellyfish Aequorea

victoria, which emits green light (emission peak at a wavelenth of 509 nm) when excited by blue light (excitation peak at a wavelenth of 395 nm). Green Fluorescent Protein (GFP) has become an invaluable tool in cell biology research, since its intrinsic fluorescence can be visualized in living cells. GFP fluorescence is stable under fixation conditions and suitable for a variety of applications. GFP has been widely used as a reporter for gene expression, enabling researchers to visualize and localize GFP-tagged proteins within living cells without the need for chemical staining. Other applications of GFP include assessment of protein protein interactions through the yeast two hybrid system and measurement of distance between proteins through fluorescence energy transfer (FRET) protocols. GFP technnology

has considerably contributed to a greater understanding of cellular physiology.

YFP differs from GFP due to a mutation at T203Y; antibodies raised against full-length GFP

should also detect YFP and other variants.

Synonyms: Green fluorescent protein, GFP-Tag