

## **Product datasheet for R1149PS**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Glucoamylase 1 Goat Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** ELISA, WB

Recommended Dilution: Western blot: 1/500-1/3,000.

**ELISA**: 1/5,000-1/25,000.

This product has been assayed against 1.0  $\mu$ g of Glucoamylase [Rhizopus] in a standard ELISA using peroxidase conjugated affinity purified anti-goat IgG (rabbit) and ABTS as a substrate for 30 minutes at room temperature. A working dilution of 1/5,000 to 1/25,000 of the

reconstitution concentration is suggested.

Reactivity: Rhizopus
Host: Goat

Clonality: Polyclonal

**Immunogen:** Glucoamylase from Rhizopus

Specificity: This antibody detects Rhizopus Glucoamylase. Cross reactivity against Glucoamylase from

other sources is unknown.

Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-goat serum as

well as purified and partially purified Glucoamylase from Rhizopus.

**Formulation:** 0.02 M Potassium phosphate, 0.15 M Sodium chloride, pH 7.2

State: Purified

State: Lyophilized purified Ig fraction Preservative: 0.01% (w/v) Sodium azide

**Reconstitution Method:** Restore with 0.1 ml of deionized water (or equivalent).

**Concentration:** lot specific

**Purification:** Delipidation, salt fractionation and ion exchange chromatography followed by extensive

dialysis against the buffer

Conjugation: Unconjugated

**Storage:** Store lyophilized at 2-8°C for 6 months or at -20°C long term.

After reconstitution store the antibody undiluted at 2-8°C for one month or (in aliquots) at -

20°C long term.

Avoid repeated freezing and thawing.





## Glucoamylase 1 Goat Polyclonal Antibody - R1149PS

**Stability:** Shelf life: one year from despatch.

Database Link: P07683

**Background:** Glucoamylase is an important industrial enzyme used in saccharification steps in both starch

enzymatic conversion and in alcohol production. The catalytic domain degrades

oligosaccharides from the non reducing end, releasing glucose, and the starch domain binds

the enzyme to raw starch and to the cell wall.

Synonyms: Gluc 1