

## Product datasheet for R1176

### Phosphotyrosine Mouse Monoclonal Antibody [Clone ID: 13F9]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	13F9
Applications:	ELISA, WB
Recommended Dilution:	<p>This monoclonal antibody reacts specifically with phosphotyrosine and shows minimal reactivity by ELISA and competitive ELISA with phosphoserine or phosphothreonine. The antibody reacts with free phosphotyrosine, phosphotyrosine conjugated to carriers such as thyroglobulin or BSA, and detects the presence of phosphotyrosine in proteins of both unstimulated and stimulated cell lysates. Although not tested, this antibody is likely functional in RIA, Flow cytometry, Immunohistochemistry and Immunoprecipitation. Phosphorylation of tyrosine residues is associated with many growth factors and oncogene protein kinases, and is important for cell signaling in activation, proliferation and differentiation.</p> <p><u>Recommended Dilutions:</u> This product was assayed by against phosphotyrosine conjugated BSA by ELISA using HRP Goat-anti-Mouse IgG for detection. A working dilution of 1:2,000 to 1:10,000 is suggested for this assay. For Immunoblotting dilute the antibody 1:1000 immediately before use.</p>
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	This monoclonal antibody was produced after repeated immunizations of balb/c mice with phosphotyrosine conjugated KLH.
Specificity:	<p>This product is clarified ascites produced in balb/c mice using clone 13F9.</p> <p>Reactivity is specific for phosphotyrosine and minimal cross reactivity is observed against phosphoserine or phosphothreonine.</p>
Formulation:	<p>PBS buffer with 0.02% sodium azide as preservative.</p> <p>State: Ascites</p> <p>State: Liquid (sterile filtered) ascites.</p>
Conjugation:	Unconjugated


[View online »](#)

- Storage:** Store the antibody (undiluted) at 2-8°C for one month or (in aliquots) at -20°C for longer. Dilute only prior to immediate use. Avoid repeated freezing and thawing.
- Stability:** Shelf life: One year from despatch.
- Note:** **Hybridoma:** Produced by the fusion between mouse splenocytes and mouse myeloma SP2/0 cells using conventional hybridoma technology.

## Product images:




Figure 1. Mab anti-phosphotyrosine antibody (clone 13F9) is shown to detect by immunoblot (lane 1) phosphorylated EGFR in an A431 cell lysate (10 g per lane) after stimulation with EGF. The binding of EGF to EGFR causes rapid activation of its intrinsic au




Figure 2. ELISA results of Mab antiphosphotyrosine antibody tested against BSA conjugates of pT, pY and pS. Each well was coated with 0.1g of conjugate. The starting dilution of antibody was 1:1000 and each point on the Y-axis represents a 2-fold dilution.