

Product datasheet for **R1188**

CREB1 (N-term) Rabbit Polyclonal Antibody

Product data:

| | |
|-----------------------|--|
| Product Type: | Primary Antibodies |
| Applications: | ELISA, EMSA, IP, WB |
| Recommended Dilution: | <p>Western Blot: 1/5,00-1/1,000.</p> <p>This antibody was assayed against a lysate of 1 x 10⁶ Raji B cells by Immunoblot and found to be reactive against CREB-1 (p43), showing a 46 kDa band at a dilution of 1/500-1/1,000 followed by reaction with Peroxidase conjugated anti-Rabbit IgG (H&L) [Goat] (Cat#R1364HRP)</p> <p>ELISA: 1/5,000-1/25,000.</p> <p>Immunoprecipitation.</p> <p>Gel Shift: This antibody was also tested in a <i>Gel Supershift Assay</i> and found to be reactive against CREB-1 (p43) using 0.5-1.0 µl per assay.</p> |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Immunogen: | CREB-1 (p43) peptide corresponding to a region near the N-terminus of the Human protein conjugated to Keyhole Limpet Hemocyanin (KLH). |
| Specificity: | <p>This product was prepared from monospecific antiserum by delipidation and Immunoabsorption against an E.coli lysate immobilized on agarose beads.</p> <p>This antibody may react non-specifically with other proteins. A partial cross-reactivity is observed against CREM-1 protein.</p> |
| Formulation: | <p>0.02M Potassium Phosphate, 0.15M Sodium Chloride, pH 7.2</p> <p>State: Liquid</p> <p>State: Liquid (sterile filtered) Serum</p> <p>Stabilizer: None</p> <p>Preservative: 0.01% (w/v) Sodium Azide</p> |
| Concentration: | lot specific |
| Conjugation: | Unconjugated |
| Storage: | <p>Store the antibody (in aliquots) at -20°C.</p> <p>Dilute only prior to immediate use.</p> <p>Avoid repeated freezing and thawing.</p> |



[View online »](#)

Stability: Shelf life: 6 month from despatch.


Gene Name: cAMP responsive element binding protein 1

Database Link: [Entrez Gene 1385 Human P16220](#)

Background: Cyclic AMP Response Element Binding protein (CREB) is a basic / leucine zipper transcription factor that binds the cyclic AMP response element (CRE) and activates transcription in response to a variety of extracellular signals including neurotransmitters, hormones, membrane depolarization, and growth and neurotrophic factors. Activation of CREB is dependent upon the phosphorylation of serine 133. Phosphorylation occurs via p44 / 42 MAP kinase and p90RSK and also via p38 MAP kinase and MSK 1. Although CREB will bind DNA independent of its phosphorylation state, only the phosphorylated form is competent as a transcription factor. CREB binding protein (CBP), a transcriptional coactivator that directly interacts with CREB, binds to CREB in the region of serine 133. CREB appears to play an important role in learning and memory. CREB knock out mice show diminished learning ability.

Synonyms: CREB-1

Product images:



CREB antibody is shown to detect CREB-1 present in Raji B cell nuclear extract lysates. Detection occurs using a 1/1,000 dilution of antibody followed by 1/5,000 dilution of HRP Goat-anti Rabbit IgG with visualization via ECL. Film exposure approximately 1'. Other detection systems will yield similar results.