

Product datasheet for **R1303F**

Chicken IgG / Chicken IgY (H+L chain), adsorbed Goat Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Chicken IgG / Chicken IgY (H+L chain), adsorbed Goat Polyclonal Antibody
Recommended Dilution:	Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. Flow cytometry: 1:2,000 - 1:10,000. IF Microscopy: 1:500 - 1:2,500.
Reactivity:	Chicken
Host:	Goat
Immunogen:	Chicken IgG whole molecule
Formulation:	PBS, pH 7.2 containing 10 mg/ml Bovine Serum Albumin (BSA; IgG and Protease free) as stabilizer and 0.01% sodium azide as a preservative Label: FITC State: Lyophilized purified Ig fraction Label: Fluorescein isothiocyanate (Molecular Weight 390 daltons) Absorption emission: 495 nm / 528 nm Molar ratio: 2.6 moles FITC per mole of Goat IgG
Reconstitution Method:	Restore with 1.0 ml of deionized water (or equivalent).
Concentration:	lot specific
Purification:	Immunoaffinity chromatography
Conjugation:	FITC
Storage:	Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid repeated freezing and thawing.



[View online »](#)