

## Product datasheet for R1303T

### Chicken IgG / Chicken IgY (H+L chain), adsorbed Goat Polyclonal Antibody

#### Product data:

<b>Product Type:</b>	Secondary Antibodies
<b>Product Name:</b>	Chicken IgG / Chicken IgY (H+L chain), adsorbed Goat Polyclonal Antibody
<b>Recommended Dilution:</b>	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency.
<b>Reactivity:</b>	Chicken
<b>Host:</b>	Goat
<b>Immunogen:</b>	Chicken IgG whole molecule
<b>Formulation:</b>	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, containing 10 mg/ml Bovine Serum Albumin (BSA) IgG and Protease free as a stabilizer and 0.01% (w/v) Sodium Azide as preservative Label: TRITC State: Lyophilized Ig fraction Label: Tetramethylrhodamine isothiocyanate (Molecular Weight 444 daltons) Absorption emission: 550 nm / 570 nm Molar ratio: 3.5 moles TRITC per mole of Goat IgG
<b>Reconstitution Method:</b>	Restore with 1.0 ml of deionized water (or equivalent)
<b>Concentration:</b>	lot specific
<b>Purification:</b>	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Chicken IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities.
<b>Conjugation:</b>	TRITC
<b>Storage:</b>	Store vial at 4°C prior to restoration. Restore with deionized water or equivalent; centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. For extended storage mix product with glycerol to 50% and then aliquot contents and freeze at -20°C or below. Avoid cycles of freezing and thawing. Dilute only prior to immediate use.



[View online »](#)