

Product datasheet for **R1321TR**

Goat IgG (F(ab)₂ specific), adsorbed Rabbit Polyclonal Antibody

Product data:

| | |
|-------------------------------|---|
| Product Type: | Secondary Antibodies |
| Product Name: | Goat IgG (F(ab) ₂ specific), adsorbed Rabbit Polyclonal Antibody |
| Recommended Dilution: | Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency. |
| Reactivity: | Goat |
| Host: | Rabbit |
| Immunogen: | Goat IgG F(ab') ₂ fragment. |
| Formulation: | 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer and 0.01% (w/v) sodium azide as preservative. Label: Texas Red State: Lyophilized purified Ig fraction. Label: (TM) Sulfonyl Chloride (Molecular Weight 625 daltons) Absorption emission: 596 nm Emission Wavelength: 620 nm Molar radio: 3.4 moles Texas Red per mole of Rabbit IgG. |
| Reconstitution Method: | Restore with 1.0 ml of deionized water (or equivalent). |
| Concentration: | lot specific |
| Purification: | Immunoaffinity chromatography. |
| Conjugation: | Texas Red |
| Storage: | Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid repeated freezing and thawing. |



[View online »](#)