

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for R1323B

Guinea pig IgG (Fc specific) Rabbit Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Guinea pig IgG (Fc specific) Rabbit Polyclonal Antibody
Recommended Dilution:	Suitable for Immunoblotting, ELISA, Immunohistochemistry, Immunomicroscopy as well as other antibody based assays using streptavidin or avidin conjugates requiring lot-to-lot consistency. <u>Note</u> : This product has been assayed against 1.0 µg of Guinea Pig IgG in a standard Capture ELISA using Peroxidase Conjugated Streptavidin and ABTS (2,2'-azino-bis-[3- ethylbenthiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1/4,000 to 1/20,000 of the reconstitution concentration is suggested for this product. <u>Recommended Dilutions</u> : ELISA: 1/20,000-1/100,000 Western Blot: 1/2,000-1/10,000 Immunohistochemistry: 1/1,000-1/5,000
Reactivity:	Guinea Pig
Host:	Rabbit
Host: Immunogen:	Rabbit Guinea Pig IgG F(c) fragment.
Immunogen:	Guinea Pig IgG F(c) fragment. 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 10 mg/ml BSA (IgG and Protease free) as stabilizer and 0.01% (w/v) sodium azide as preservative. Label: Biotin State: Lyophilized purified Ig fraction.
Immunogen: Formulation:	Guinea Pig IgG F(c) fragment. 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 10 mg/ml BSA (IgG and Protease free) as stabilizer and 0.01% (w/v) sodium azide as preservative. Label: Biotin State: Lyophilized purified Ig fraction. Molar radio: 10-20 BAC molecules per Goat IgG molecule.
Immunogen: Formulation: Reconstitution Method:	Guinea Pig IgG F(c) fragment. 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 10 mg/ml BSA (IgG and Protease free) as stabilizer and 0.01% (w/v) sodium azide as preservative. Label: Biotin State: Lyophilized purified Ig fraction. Molar radio: 10-20 BAC molecules per Goat IgG molecule. Restore with 1.0 ml of deionized water (or equivalent).
Immunogen: Formulation: Reconstitution Method: Concentration:	Guinea Pig IgG F(c) fragment. 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 10 mg/ml BSA (IgG and Protease free) as stabilizer and 0.01% (w/v) sodium azide as preservative. Label: Biotin State: Lyophilized purified Ig fraction. Molar radio: 10-20 BAC molecules per Goat IgG molecule. Restore with 1.0 ml of deionized water (or equivalent). 2.0 mg/ml (by UV absorbance at 280 nm)



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Guinea pig lgG (Fc specific) Rabbit Polyclonal Antibody – R1323B
Storage:	Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid repeated freezing and thawing.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US