

## Product datasheet for R1327B

### Hamster IgG (H+L chain) Rabbit Polyclonal Antibody

#### Product data:

<b>Product Type:</b>	Secondary Antibodies
<b>Product Name:</b>	Hamster IgG (H+L chain) Rabbit Polyclonal Antibody
<b>Recommended Dilution:</b>	Suitable for immunoblotting, ELISA, immunohistochemistry, immunomicroscopy as well as other antibody based assays using streptavidin or avidin conjugates requiring lot-to-lot consistency. This product has been assayed against 1.0µg of hamster IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin and ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1/15,000 to 1/60,000 of the reconstitution concentration is suggested for this product.
<b>Reactivity:</b>	Hamster
<b>Host:</b>	Rabbit
<b>Immunogen:</b>	Hamster IgG whole molecule
<b>Formulation:</b>	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, containing 10 mg/ml Bovine Serum Albumin (BSA) IgG and Protease free, and 0.01% (w/v) Sodium Azide Label: Biotin State: Lyophilized IgG fraction Label: Biotinamidocaproate N-Hydroxysuccinimide Ester (BAC) Molar ratio: 10-20 BAC molecules per Rabbit IgG molecule
<b>Reconstitution Method:</b>	Restore with 1.0 ml of deionized water or equivalent.
<b>Concentration:</b>	lot specific
<b>Purification:</b>	Immunoaffinity chromatography using Hamster IgG coupled to agarose
<b>Conjugation:</b>	Biotin
<b>Storage:</b>	Store vial at 4°C prior to restoration. Restore with deionized water (or equivalent); centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4°C as an undiluted liquid. For extended storage reconstitute product with 50% glycerol instead of water and then aliquot contents and freeze at -20°C or below. Avoid cycles of freezing and thawing. Dilute only prior to immediate use.



[View online »](#)