

Product datasheet for **R1330TR**

Equine IgG (Fc specific) Rabbit Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Equine IgG (Fc specific) Rabbit Polyclonal Antibody
Recommended Dilution:	Suitable for immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency.
Reactivity:	Horse
Host:	Rabbit
Immunogen:	Horse IgG F(c) fragment
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2; 10 mg/ml Bovine Serum Albumin (BSA) IgG and Protease free; 0.01% (w/v) Sodium Azide Label: Texas Red State: Lyophilized Ig fraction Label: Texas Red™ Sulfonyl Chloride (Molecular Weight 625 daltons) Absorption emission: 596 nm / 620 nm Molar ratio: 2.0 moles Texas Red™ per mole of Rabbit IgG
Reconstitution Method:	Restore with 1.0 ml of deionized water (or equivalent).
Concentration:	lot specific
Purification:	Prepared from monospecific antiserum by immunoaffinity chromatography using Horse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities
Conjugation:	Texas Red
Storage:	Store vial at 2-8 ° C prior to restoration. Following restoration product can be stored undiluted at 2-8 ° for up to one month or (in aliquots) at -20 ° C or below. Avoid repeated freezing and thawing. Centrifuge product if not completely clear after standing at room temperature.



[View online »](#)