

Product datasheet for **R1340B**

Human IgM (Mu chain specific) Goat Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Human IgM (Mu chain specific) Goat Polyclonal Antibody
Recommended Dilution:	Suitable for Immunoblotting (1:2,000 - 1:10,000), ELISA (1:20,000 - 1:100,000), Immunohistochemistry (1:1,000 - 1,5,000), Immunomicroscopy as well as other antibody based assays using streptavidin or avidin conjugates requiring lot-to-lot consistency. Recommended Dilutions: This product has been assayed against 1.0 µg of Human IgM in a standard capture ELISA using Peroxidase Conjugated Streptavidin and ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:3,000 to 1:17,000 of the reconstitution concentration is suggested for this product.
Reactivity:	Human
Host:	Goat
Immunogen:	Human IgM µ chain.
Formulation:	0.01 M Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.4 with 10 mg/ml Bovine Serum Albumin (BSA) IgG and Protease free as stabilizer and 0.01% (w/v) Sodium Azide as preservative. Label: Biotin State: Lyophilized purified Ig fraction. Label: Biotinamidocaproate N-Hydroxysuccinimide Ester (BAC) Molar radio: 10-20 BAC molecules per Goat IgG molecule.
Reconstitution Method:	Restore with 1.0 ml of deionized water (or equivalent).
Concentration:	lot specific
Purification:	Immunoaffinity chromatography.
Conjugation:	Biotin
Storage:	Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid repeated freezing and thawing.



[View online »](#)