

## Product datasheet for R1347T

### Mouse IgG (H+L chain) Chicken Polyclonal Antibody

#### Product data:

<b>Product Type:</b>	Secondary Antibodies
<b>Product Name:</b>	Mouse IgG (H+L chain) Chicken Polyclonal Antibody
<b>Applications:</b>	FC, IF
<b>Recommended Dilution:</b>	Immunomicroscopy. Flow Cytometry or FACS analysis.
<b>Reactivity:</b>	Mouse
<b>Host:</b>	Chicken
<b>Immunogen:</b>	Mouse IgG whole molecule.
<b>Formulation:</b>	0,02 M Potassium Phosphate, 0,15 M Sodium Chloride, pH 7,2 containing 10 mg/ml BSA (IgG and Protease free) as stabilizer and 0,01% (w/v) Sodium Azide as preservative. Label: TRITC State: Lyophilized purified Ig fraction. Label: Tetramethylrhodamine isothiocyanate (MW 444 daltons) Absorption emission: 550 nm / 570 nm Molar ratio: 3.0 moles TRITC per mol of Chicken IgG
<b>Reconstitution Method:</b>	Restore with 1.0 ml of deionized water (or equivalent)
<b>Concentration:</b>	lot specific
<b>Purification:</b>	Prepared from monospecific antiserum by Immunoaffinity Chromatography using Mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities.
<b>Conjugation:</b>	TRITC
<b>Storage:</b>	Store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.



[View online »](#)