

Product datasheet for R1365T

Rabbit IgG (H+L chain, adsorbed) Goat Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Rabbit IgG (H+L chain, adsorbed) Goat Polyclonal Antibody
Recommended Dilution:	Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency.
Reactivity:	Rabbit
Host:	Goat
Immunogen:	Rabbit IgG whole molecule.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 containing 10 mg/ml Bovine Serum Albumin (BSA) IgG and Protease free and 0.01% (w/v) Sodium Azide conjugated to Label: TRITC State: Lyophilized purified Ig fraction Label: Tetramethylrhodamine isothiocyanate (Molecular Weight 444 daltons) Absorption emission: 550 nm / 570 nm Molar ratio: 3.2 moles TRITC per mole of Goat IgG.
Reconstitution Method:	Restore with 1.0 ml of deionized water (or equivalent).
Concentration:	lot specific
Purification:	Immunoaffinity chromatography.
Conjugation:	TRITC
Storage:	Store vial at 2-8°C prior to restoration. For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid repeated freezing and thawing.



[View online »](#)