

Product datasheet for **R1378B**

Rat IgG (H+L chain, adsorbed) Goat Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Rat IgG (H+L chain, adsorbed) Goat Polyclonal Antibody
Recommended Dilution:	Suitable for Immunoblotting (1/2,000-1/10,000), ELISA, Immunohistochemistry (1/1,000-1/5,000), Immunomicroscopy as well as other antibody based assays using streptavidin or avidin conjugates requiring lot-to-lot consistency. Recommended Dilutions: This product has been assayed against 1.0 µg of Rat IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin and ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:20,000 to 1:100,000 of the reconstitution concentration is suggested for this product.
Reactivity:	Rat
Host:	Goat
Immunogen:	Rat IgG whole molecules.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 containing 10 mg/ml Bovine Serum Albumin (BSA IgG and Protease free) and 0.01% (w/v) Sodium Azide as preservative. Label: Biotin State: Lyophilized purified Ig fraction. Label: Biotinamidocaproate N-Hydroxysuccinimide Ester (BAC) Molar ratio: 10-20 BAC molecules per Goat IgG molecule.
Reconstitution Method:	Restore with 1.0 ml of deionized water (or equivalent).
Concentration:	lot specific
Purification:	Immunoaffinity chromatography.
Conjugation:	Biotin
Storage:	Store vial at 4°C prior to restoration. For extended storage reconstitute product with 50% glycerol instead of water and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 4°C as an undiluted liquid. Dilute only prior to immediate use. Avoid cycles of freezing and thawing.



[View online »](#)