

## Product datasheet for R1387F

### Goat IgG (H+L chain), Fab Fragment, adsorbed Donkey Polyclonal Antibody

#### Product data:

<b>Product Type:</b>	Secondary Antibodies
<b>Product Name:</b>	Goat IgG (H+L chain), Fab Fragment, adsorbed Donkey Polyclonal Antibody
<b>Recommended Dilution:</b>	Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. Recommended dilutions: Flow cytometry 1:2,000 - 1:10,000 Immunofluorescence microscopy 1:500 - 1:2,500.
<b>Reactivity:</b>	Goat
<b>Host:</b>	Donkey
<b>Immunogen:</b>	Goat IgG whole molecule.
<b>Formulation:</b>	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2, containing 10 mg/ml Bovine Serum Albumin (BSA) IgG and Protease free, and 0.01% (w/v) Sodium Azide as preservative. Label: FITC State: Lyophilized Fab fragments Label: Fluorescein isothiocyanate (Molecular Weight 390 daltons) Absorption emission: 495 nm / 528 nm Molar ratio: 1.7 moles FITC per mole of Donkey IgG Fab.
<b>Reconstitution Method:</b>	Restore with 1.0 ml of deionized water or equivalent.
<b>Concentration:</b>	lot specific
<b>Purification:</b>	Immunoaffinity chromatography.
<b>Conjugation:</b>	FITC
<b>Storage:</b>	Store vial at 4°C prior to restoration. centrifuge product if not completely clear after standing at room temperature. This product is stable for one month at 4°C as an undiluted liquid. For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Avoid cycles of freezing and thawing. Dilute only prior to immediate use.



[View online »](#)