

Product datasheet for **R1388B**

Human IgG (H+L chain), Fab Fragment Goat Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Human IgG (H+L chain), Fab Fragment Goat Polyclonal Antibody
Recommended Dilution:	Suitable for Immunoblotting, ELISA, Immunohistochemistry, Immunomicroscopy as well as other antibody based assays using streptavidin or avidin conjugates requiring lot-to-lot consistency. <u>Recommended Dilutions:</u> This product has been assayed against 1.0 ug of Human IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin and ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6- sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1/10,000-1/50,000 of the reconstitution concentration is suggested for this product.
Reactivity:	Human
Host:	Goat
Immunogen:	Human IgG whole molecule.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 10 mg/ml Bovine Serum Albumin (BSA) IgG and Protease free as stabilizer and 0.01% (w/v) Sodium Azide as preservative. Label: Biotin State: Lyophilized Fab fragments. Label: Biotinamidocaproate N-Hydroxysuccinimide Ester (BAC) Molar radio: 10-20 BAC molecules per Goat IgG Fab molecule.
Reconstitution Method:	Restore with 1.0 ml of deionized water (or equivalent).
Concentration:	lot specific
Purification:	Immunoaffinity Chromatography.
Conjugation:	Biotin
Storage:	Store vial at 4°C prior to restoration. For extended storage reconstitute product with 50% glycerol instead of water and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid cycles of freezing and thawing.



[View online »](#)