

## Product datasheet for R1392T

### Goat IgG (H+L chain), F(ab)<sub>2</sub> Fragment, adsorbed Rabbit Polyclonal Antibody

#### Product data:

<b>Product Type:</b>	Secondary Antibodies
<b>Product Name:</b>	Goat IgG (H+L chain), F(ab) <sub>2</sub> Fragment, adsorbed Rabbit Polyclonal Antibody
<b>Recommended Dilution:</b>	Suitable for Immunomicroscopy and Flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring lot-to-lot consistency.
<b>Reactivity:</b>	Goat
<b>Host:</b>	Rabbit
<b>Immunogen:</b>	Goat IgG whole molecule.
<b>Formulation:</b>	0.01 M Sodium Phosphate, 0.25 M Sodium Chloride, pH 7.6 containing 15 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer and 0.01% (w/v) Sodium Azide, 0.01% (w/v) Thimerosal as preservative. Label: TRITC State: Lyophilized F(ab') <sub>2</sub> fragments. Label: Tetramethylrhodamine isothiocyanate ; Molecular Weight 444 daltons Absorption emission: 550 nm / 570 nm Molar ratio: 1.6 moles TRITC per mole of Rabbit IgG Fab).
<b>Reconstitution Method:</b>	Restore with 1.0 ml of deionized water (or equivalent).
<b>Concentration:</b>	lot specific
<b>Purification:</b>	Immunoaffinity chromatography.
<b>Conjugation:</b>	TRITC
<b>Storage:</b>	Store vial at 4°C prior to restoration. For extended storage reconstitute product with 50% glycerol instead of water and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 4°C as an undiluted liquid. Dilute only prior to immediate use. Avoid cycles of freezing and thawing.



[View online »](#)