

Product datasheet for **R1396AP**

Human IgG (adsorbed F(ab)2 Fragment) Goat Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Human IgG (adsorbed F(ab)2 Fragment) Goat Polyclonal Antibody
Recommended Dilution:	Suitable for Immunoblotting (Western or Dot blot), ELISA and Immunohistochemistry as well as other phosphatase-antibody based enzymatic assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. This product has been assayed against 1.0 ug of Human IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate as a substrate for 30 minutes at room temperature. A working dilution of 1:5,000 to 1:20,000 is suggested for this product.
Reactivity:	Human
Host:	Goat
Immunogen:	Human IgG whole molecule.
Formulation:	0.01 M Tris Chloride, 0.15M Sodium Chloride, 0.001 M Magnesium Chloride, 0.0001 M Zinc Chloride, 50% (v/v) Glycerol; pH 8.0 with 10 mg/ml Bovine Serum Albumin (BSA) IgG and Protease free as stabilizer and 0.05% (w/v) sodium azide as preservative. Label: AP State: Liquid (sterile filtered) F(ab')2 fragments. Label: Alkaline Phosphatase (Calf Intestine) (Molecular Weight 140,000 daltons)
Concentration:	lot specific
Purification:	Immunoaffinity chromatography.
Conjugation:	AP
Storage:	Store the antibody as an undiluted liquid at 4°C. DO NOT FREEZE! Dilute only prior to immediate use. Freezing alkaline phosphatase conjugates will result in a substantial loss of enzymatic activity. Dilute only prior to immediate use.



[View online »](#)