

## Product datasheet for R1399AP

### Human IgA + IgG + IgM (adsorbed F(ab)2 Fragment) Goat Polyclonal Antibody

#### Product data:

Product Type:	Secondary Antibodies
Product Name:	Human IgA + IgG + IgM (adsorbed F(ab)2 Fragment) Goat Polyclonal Antibody
Recommended Dilution:	Suitable for Immunoblotting (Western or dot blot), ELISA and immunohistochemistry as well as other phosphatase-antibody based enzymatic assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. This product has been assayed against 1.0 ug of Human IgG in a standard capture ELISA using pNPP p-nitrophenyl phosphate as a substrate for 30 minutes at room temperature. A working dilution of 1:1,000 to 1:5,000 of the reconstitution concentration is suggested for this product.
Reactivity:	Human
Host:	Goat
Immunogen:	Human IgG whole molecule
Formulation:	0.01M Tris Chloride, 0.15M Sodium Chloride, 0.001M Magnesium Chloride, 0.0001M Zinc Chloride, 50% (v/v) Glycerol; pH 8.0 with 10 mg/ml BSA (IgG and Protease free) as stabilizer and 0.01% (w/v) sodium azide as preservative. Label: AP State: Liquid (sterile filtered) F(ab')2 fragments Label: Alkaline Phosphatase (Calf Intestine) (Molecular Weight 140,000 daltons)
Concentration:	lot specific
Purification:	Immunoaffinity chromatography
Conjugation:	AP
Storage:	This antibody is stable at 2-8°C as an undiluted liquid. <b>DO NOT FREEZE!</b> Store reagent in the dark. Use subdued lighting during handling and incubation of cells prior to analysis. Dilute only prior to immediate use.



[View online »](#)