

Product datasheet for R1407B

Mouse IgG (F(ab)2 specific), F(ab)2 Fragment, adsorbed Goat Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Mouse IgG (F(ab)2 specific), F(ab)2 Fragment, adsorbed Goat Polyclonal Antibody
Recommended Dilution:	Suitable for Immunoblotting, ELISA, Immunohistochemistry, immunomicroscopy as well as other antibody based assays using streptavidin or avidin conjugates requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. <u>Recommended Dilutions:</u> This product has been assayed against 1.0 ug of Mouse IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin and ABTS (2,2'-azino-bis-[3-ethylbenzothiazoline-6- sulfonic acid]) as a substrate for 30 minutes at room temperature. A working dilution of 1:4,000 to 1:20,000 of the reconstitution concentration is suggested for this product.
Reactivity:	Mouse
Host:	Goat
Immunogen:	Mouse IgG F(ab')2 fragment
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 10 mg/ml Bovine Serum Albumin (BSA) IgG and Protease free as stabilizer and 0.01% (w/v) Sodium Azide as preservative. Label: Biotin State: Lyophilized F(ab')2 fragments. Label: Biotinamidocaproate N-Hydroxysuccinimide Ester (BAC) Molar ratio: 10-20 BAC molecules per Goat IgG Fab molecule.
Reconstitution Method:	Restore with 1.0 ml of deionized water (or equivalent).
Concentration:	lot specific
Purification:	Immunoaffinity chromatography.
Conjugation:	Biotin



[View online »](#)

Storage:

Store vial at 2-8°C prior to restoration. Restore with 1.0 ml of deionized water (or equivalent). For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid repeated freezing and thawing.