

Product datasheet for R1409F

Mouse IgA (H+L chain), F(ab)₂ Fragment, adsorbed Goat Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Mouse IgA (H+L chain), F(ab) ₂ Fragment, adsorbed Goat Polyclonal Antibody
Recommended Dilution:	Suitable for Immunomicroscopy and Flow Cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.
Reactivity:	Mouse
Host:	Goat
Immunogen:	Mouse IgG, IgA, and IgM whole molecules.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 0.01% (w/v) Sodium Azide as preservative and 10 mg/ml Bovine Serum Albumin (BSA) as stabilizer. Label: FITC State: Lyophilized F(ab') ₂ fragments. Label: Fluorescein isothiocyanate (Molecular Weight 390 daltons) Absorption emission: 495 nm / 528 nm Molar ratio: 4.0 moles FITC per mole of Goat IgG F(ab') ₂ .
Reconstitution Method:	Restore with 1.0 ml of deionized water (or equivalent).
Concentration:	lot specific
Purification:	Immunoaffinity chromatography.
Conjugation:	FITC
Storage:	Store vial at 2-8°C prior to restoration. Restore with 1.0 ml of deionized water (or equivalent). For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid repeated freezing and thawing.



[View online »](#)