

## Product datasheet for R1412T

### Rabbit IgG (F(ab)2 specific adsorbed F(ab)2 Fragment) Goat Polyclonal Antibody

#### Product data:

Product Type:	Secondary Antibodies
Product Name:	Rabbit IgG (F(ab)2 specific adsorbed F(ab)2 Fragment) Goat Polyclonal Antibody
Recommended Dilution:	Suitable for Immunomicroscopy and flow cytometry or FACS analysis as well as other antibody based fluorescent assays requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity.
Reactivity:	Rabbit
Host:	Goat
Immunogen:	Rabbit IgG F(ab')2 fragment.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 containing 10 mg/ml Bovine Serum Albumin (BSA, IgG and Protease free) as stabilizer and 0.01% (w/v) Sodium Azide as preservative. Label: TRITC State: Lyophilized F(ab')2 fragments. Label: Tetramethylrhodamine isothiocyanate ; Molecular Weight 444 daltons) Absorption emission: 550 nm / 570 nm Molar ratio: 1.4 moles TRITC per mole of Goat IgG F(ab')2).
Reconstitution Method:	Restore with 0.5 ml of deionized water (or equivalent).
Concentration:	lot specific
Purification:	Immunoaffinity chromatography.
Conjugation:	TRITC
Storage:	Store the antibody at 2-8°C prior to restoration. Restore with 0.5 ml of deionized water (or equivalent). For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid repeated freezing and thawing.



[View online »](#)