

Product datasheet for R1413B

Rat IgG (H+L chain), F(ab)2 Fragment Rabbit Polyclonal Antibody

Product data:

Product Type:	Secondary Antibodies
Product Name:	Rat IgG (H+L chain), F(ab)2 Fragment Rabbit Polyclonal Antibody
Recommended Dilution:	Suitable for Immunoblotting, ELISA, Immunohistochemistry, Immunomicroscopy as well as other antibody based assays using streptavidin or avidin conjugates requiring extremely low background levels, absence of F(c) mediated binding, lot-to-lot consistency, high titer and specificity. <u>Recommended Dilution(s)</u> : This product has been assayed against 1.0 µg of Rat IgG in a standard capture ELISA using Peroxidase Conjugated Streptavidin and ABTS as a substrate for 30 minutes at room temperature. A working dilution of 1:10,000 to 1:50,000 of the reconstitution concentration is suggested.
Reactivity:	Rat
Host:	Rabbit
Immunogen:	Rat IgG whole molecule
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 containing 10 mg/ml BSA (IgG and Protease free) as stabilizer and 0.01% (w/v) Sodium Azide preservative . Label: Biotin State: Lyophilized F(ab')2 fragments. Label: Biotinamidocaproate N-Hydroxysuccinimide Ester (BAC) Molar ratio: 10-20 BAC molecules per Rabbit IgG molecule.
Reconstitution Method:	Restore with 0.5 ml of deionized water (or equivalent).
Concentration:	lot specific
Purification:	Immunoaffinity chromatography.
Conjugation:	Biotin
Storage:	Store vial at 2-8°C prior to restoration. Restore with 0.5 ml of deionized water (or equivalent). For extended storage add glycerol to 50% and then aliquot contents and freeze at -20°C or below. Centrifuge product if not completely clear after standing at room temperature. This antibody is stable for one month at 2-8°C as an undiluted liquid. Dilute only prior to immediate use. Avoid repeated freezing and thawing.



[View online »](#)