

Product datasheet for **R1564P**

ULP1 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	This purified polyclonal antibody reacts with yeast ULP-1 by Western blot (1/500-1:2,000) and ELISA (1/4,000-1/20,000). Although not tested, this antibody is likely functional in Immunohistochemistry and Immunoprecipitation. Expect a band approximately 72.4 kDa in size corresponding to yeast ULP-1 by western blotting in the appropriate lysate or extract.
Reactivity:	Yeast
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Prepared from rabbit serum after repeated immunizations with recombinant yeast ULP-1 protein.
Specificity:	Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum. Reactivity against ULP-1 from other sources or ULP-2 has not been determined.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 with 0.01% (w/v) Sodium Azide as preservative. State: Purified State: Lyophilized purified Ig fraction.
Reconstitution Method:	Restore with 0.1 ml of deionized water (or equivalent).
Concentration:	lot specific
Purification:	Multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis.
Conjugation:	Unconjugated
Storage:	Store vial at 2-8°C prior to restoration. After restoration, store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing. Centrifuge product if not completely clear after standing at room temperature.
Stability:	Shelf life: one year from despatch.



[View online »](#)

Database Link: [Q02724](#)

Background: ULP-1, ubiquitin-like protein-specific protease 1, initially processes Smt3 and also acts as a deconjugating enzyme for Smt3 [*Saccharomyces cerevisiae* (Baker's yeast)]. Covalent modification of cellular proteins by the ubiquitin-like modifier SUMO (small ubiquitin-like modifier) regulates various cellular processes, such as nuclear transport, signal transduction, stress responses and cell cycle progression. But, in contrast to ubiquitination, sumoylation does not tag proteins for degradation by the 26S proteasome, but rather seems to enhance stability or modulate their subcellular compartmentalization. Once covalently attached to cellular targets, SUMO regulates protein:protein and protein:DNA interactions, as well as localization and stability of the target protein. Sumoylation occurs in most eukaryotic systems, and SUMO is highly conserved from yeast to humans. Where invertebrates have only a single SUMO gene termed SMT3, three members of the SUMO family have been identified in vertebrates: SUMO-1 and the close homologues SUMO-2 and SUMO-3. Three distinct steps can be distinguished in the SUMO modification pathway: 1) activation of SUMO, 2) transfer of SUMO to the conjugating enzyme, and 3) substrate modification. Since SUMO is synthesized as a precursor protein, a maturation step precedes the activation reaction. In yeast, C-terminal processing of the SUMO precursor is mediated by the processing protease Ulp1, which has an additional role in the deconjugation of SUMO-modified substrates. Mature SUMO is activated by SUMO-activating enzyme, an E1-like heterodimeric protein complex composed of Uba2 and Aos1. Ulp1 function has provided evidence that SUMO modification in yeast, as has been suspected for vertebrates, plays an important role in nucleocytoplasmic trafficking.

Synonyms: Ubiquitin-like-specific protease 1, LPB11C, YPL020C

Note: **Protein Sequence :** Yeast ULP-1, 621 aa, predicted MW 72.4 kDa
1 msvevdkhrn tlqyhkknp spfspisty rcyprvlnnp sesrrsasfs giykkrtnts
61 rfnylndrrv lsmeesmkdg sdraskagfi ggiretlwns gkylwhtfvk neprnfdgse
121 veasgnsdve srssgsrstd vpyglrenys sdtrkhkfdt stwalpnkrr riesegvgtp
181 stspisslas qksncdsdns itfsrdpfgw nkwktsaigs nsenntsdqk nsydrdqygt
241 afirkkkvak qninntklvs raqseevtyl rqifngeykv pkilkeerer qlklmdmdke
301 kdtglkksii dltekiktil iennknrlqt rnendddlvf vkekissle rkhkdylnqk
361 lkfdrsilef ekdfkrynei lnerkkiqed lkkkkeqlak kklvpelnek dddqvqkala
421 srentqlmnr dneitvrdv ktlaprrwln dtiieffmky iekstptntva fnsffytlns
481 ergyqgvrrw mkrkkqtqdk ldkiptpinl nqshwalgii dlkkktigyv dslsngpnam
541 sfailtdlqk yvmeeskhti gedfdlihd cpqqpnydc giyvcmntly gsadapldfd
601 ykdairmrrf iahlildal k

Product images:




Figure 1. Western blot using Affinity Purified anti-Yeast ULP-1 antibody shows detection of a truncated ULP-1 fusion protein (arrowhead). Increasing concentrations of yeast ULP-1 were run on a SDS-PAGE, transferred onto nitrocellulose, and blocked for 1 hour with 5% non-fat dry milk in TTBS, and probed overnight at 4°C with a 1:1000 dilution of anti-yULP-1 antibody in 5% non-fat dry milk in TTBS. Detection occurred using a 1:1,000 dilution of HRP-labeled Donkey anti-Rabbit IgG for 1 hour at room temperature. A chemiluminescence system was used for signal detection (Roche) using a 3-sec exposure time.




Figure 2. Western blot using Affinity Purified anti-Yeast ULP-1 antibody was used to confirm the specificity of the antibody. SDS-PAGE of 2 µg of ULP-1 homologues from other sources (lanes 2 through 9). After blocking for 1 hour with 5% non-fat dry milk in TTBS, the blot was probed overnight at 4°C with a 1:1,000 dilution of antiyULP1 antibody detected as above. This antibody is specific for yeast ULP1 and does not react with ULP1 from related sources including human SENP.