

Product datasheet for **R1619B**

Albumin (ALB) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, IHC, WB
Recommended Dilution:	Western Blot: 1/1,000 - 1/5,000. ELISA: 1/170,000. Immunohistochemistry: 1/500 - 1/3,000.
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Human Serum Albumin
Specificity:	This antibody reacts to Serum Albumin.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 containing 10 mg/ml Bovine Serum Albumin (BSA) as stabilizer and 0.01% (w/v) Sodium Azide as preservative Label: Biotin State: Lyophilized Label: Biotinamidocaproate N-Hydroxysuccinimide Ester (BAC) Molar ratio: 10-20 BAC molecules per Rabbit IgG molecule
Reconstitution Method:	Restore with 1.0 ml of deionized water (or equivalent). For extended storage add glycerol to 50%
Concentration:	lot specific
Purification:	Affinity chromatography on Protein A
Conjugation:	Biotin
Storage:	Prior to reconstitution store at 2-8°C. Following reconstitution store the antibody undiluted at 2-8°C for one month or (in aliquots) at -20°C for longer. Avoid repeated freezing and thawing.
Stability:	Shelf life: one year from despatch.
Gene Name:	albumin



[View online »](#)

Database Link: [Entrez Gene 213 Human P02768](#)

Background: Albumin is a soluble, monomeric protein which comprises about one half of the blood serum protein. Albumin functions primarily as a carrier protein for steroids, fatty acids, and thyroid hormones and plays a role in stabilizing extracellular fluid volume. Mutations in this gene on chromosome 4 result in various anomalous proteins. Albumin is a globular unglycosylated serum protein of molecular weight 65,000. The human albumin gene is 16,961 nucleotides long from the putative 'cap' site to the first poly(A) addition site. It is split into 15 exons which are symmetrically placed within the 3 domains that are thought to have arisen by triplication of a single primordial domain. Albumin is synthesized in the liver as preproalbumin which has an N terminal peptide that is removed before the nascent protein is released from the rough endoplasmic reticulum. The product, proalbumin, is in turn cleaved in the Golgi vesicles to produce the secreted albumin.

Synonyms: ALB, BSA, HSA, Serum Albumin