

Product datasheet for RA109

Goat IgG (Fc Fragment) Goat Protein

Product data:

Product Type:	Native Proteins
Description:	Goat IgG (Fc Fragment) goat protein, 1 mg
Species:	Goat
Protein Source:	Serum
Concentration:	lot specific
Purity:	This product was prepared from normal serum delipidation, salt fractionation, ion exchange chromatography followed by papain digestion and extensive dialysis against the buffer stated above.
Conjugation:	Biotin
Buffer:	State: Lyophilized purified Ig fraction. Buffer System: 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 containing 10 mg/ml Bovine Serum Albumin (BSA; IgG and Protease free) as stabilizer and 0.01% (w/v) Sodium Azide as preservative. Label: Biotinamidocaproate N-Hydroxysuccinimide Ester (BAC) Biotin/Protein Ratio: 10-20 BAC molecules per Goat IgG F(c) molecule Presentation Label: Biotin
Reconstitution Method:	Restore with 1.0 ml of deionized water (or equivalent).
Preparation:	Lyophilized purified Ig fraction.
Applications:	Immunoblotting, ELISA, Immunohistochemistry, Immunomicroscopy as well as other antibody based assays using streptavidin or avidin conjugates. Assay by Immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat IgG, anti-Goat IgG F(c) and anti-Goat Serum. No reaction was observed against anti-Goat IgG F(ab') ₂ or anti-Papain.
Protein Description:	Biotin Conjugated Goat IgG F(c) fragment
Storage:	Store (undiluted) at 2-8°C prior to restoration. For extended storage reconstitute product with 50% glycerol instead of water and then aliquot contents and freeze at -20°C or below. Avoid cycles of freezing and thawing.



[View online »](#)

Stability:

Shelf life: one year from despatch.
Centrifuge product if not completely clear after standing at room temperature.
Dilute only prior to immediate use.