

Product datasheet for RA22101-100

Gfap Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Formulation:	State: Serum
Gene Name:	Mus musculus glial fibrillary acidic protein (Gfap), transcript variant 2
Database Link:	Entrez Gene 24387 RatEntrez Gene 14580 Mouse
Synonyms:	Glial Fibrillary Acidic Protein

Product images:



Mixed neuron-glial cultures stained with rabbit GFAP (red channel) and chicken vimentin (green channel). The fibroblastic cells contain only vimentin and so are green, while astrocytes contain either vimentin and GFAP, so appearing golden, or predominantly GFAP, in which case they appear red. Blue is nuclear DNA stain. Protocol on data-sheet



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2020 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn



Injection of rhHsp70 did not result in overt changes in glial cell activation. Lumbar spinal cord tissue was collected from treated and untreated G93A SOD1 or WT mice on P90. Twenty-five micrograms of protein per lane was loaded into each lane and Western blotting was performed to assess relative levels of GFAP and Iba1. When normalized to actin, quantification of the bands confirmed no change in expression (data not shown). doi:10.1523/JNEUROSCI.4057-07.2007



G93A

G93A WT WT

50 KDa

17 KDa

42 KDa

Strain:

GFAP

Iba1

Actin

Injection: rhHsp70

Mixed neuron-glial cultures stained with rabbit GFAP (red channel) and chicken vimentin (green channel). The fibroblastic cells contain only vimentin and so are green, while astrocytes contain either vimentin and GFAP, so appearing golden, or predominantly GFAP, in which case they appear red. Blue is nuclear DNA stain. Protocol on data-sheet

Injection of rhHsp70 did not result in overt changes in glial cell activation. Lumbar spinal cord tissue was collected from treated and untreated G93A SOD1 or WT mice on P90. Twenty-five micrograms of protein per lane was loaded into each lane and Western blotting was performed to assess relative levels of GFAP and Iba1. When normalized to actin, quantification of the bands confirmed no change in expression (data not shown). doi:10.1523/JNEUROSCI.4057-07.2007

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2020 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US