

## **Product datasheet for RC200014**

## MRPS23 (NM 016070) Human Tagged ORF Clone

## **Product data:**

**Product Type:** Expression Plasmids

Tag: Myc-DDK Symbol: MRPS23

Synonyms: CGI-138; COXPD46; HSPC329; MRP-S23

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)ORF Nucleotide>RC200014 ORF sequence

Sequence: Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGGCAGGCAGCCGGCTGGAAACCGTAGGGAGCATCTTCTCTCGGACTCGGGACCTGGTTCGGGCCGGGG TGCTGAAGGAGAAGCCCCTGTGGTTTGACGTATATGACGCCTTTCCCCCGCTGAGGGAGCCCGTCTTCCA AAGGCCTCGAGTGCGATATGGCAAAGCCAAAGCTCCCATCCAAGACATCTGGTACCACGAGGATCGGATT AGAGCGAAGTTTTATTCAGTGTATGGGTCTGGTCAAAGAGCTTTTGATCTATTCAATCCAAACTTCAAGT CTACCTGTCAACGGTTTGTGGAGAAGTACACTGAGCTACAGAAACTTGGAGAAACAGATGAAGAAGATT ATTTGTGGAAACAGGGAAGGCTTTATTGGCAGAAAGGTGCATTTTAAGACGAGAACCAGACGAAGCAC CAACACGGAGGTAGTCACGTTTCCCGGAAATCCGAACACTTGAGTGTCAGACCACAGACTGCGTTGGAAG AAACAGAGACTCAGAAAGAAGTTCCACAGGACCAGCATTTGGAGGCACCTGCAGACCAGTCGAAAGGTCT

CTTGCCTCCC

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC200014 protein sequence

Red=Cloning site Green=Tags(s)

MAGSRLETVGSIFSRTRDLVRAGVLKEKPLWFDVYDAFPPLREPVFQRPRVRYGKAKAPIQDIWYHEDRI RAKFYSVYGSGQRAFDLFNPNFKSTCQRFVEKYTELQKLGETDEEKLFVETGKALLAEGVILRRVGEART

QHGGSHVSRKSEHLSVRPQTALEENETQKEVPQDQHLEAPADQSKGLLPP

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: <a href="https://cdn.origene.com/chromatograms/mk6388">https://cdn.origene.com/chromatograms/mk6388</a> f12.zip




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200


CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com ORIGENE

**Restriction Sites:** Sgfl-Mlul

**Cloning Scheme:** 





<sup>\*</sup> The last codon before the Stop codon of the ORF

**ACCN:** NM\_016070

ORF Size: 570 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**Note:** Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

RefSeq: <u>NM 016070.4</u>

RefSeq Size: 1498 bp RefSeq ORF: 573 bp



**Locus ID:** 51649

 UniProt ID:
 Q9Y3D9

 Cytogenetics:
 17q22

MW: 21.8 kDa

Gene Summary: Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in


protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that

the latter contain a 5S rRNA. Among different species, the proteins comprising the


mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 28S subunit protein. A pseudogene corresponding to this gene is found on chromosome 7p. [provided by RefSeq, Jul

2008]

## **Product images:**



Circular map for RC200014



Western blot validation of overexpression lysate (Cat# [LY414209]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC200014 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).