

## Product datasheet for **RC200022L2V**

### Antizyme inhibitor 1 (AZIN1) (NM\_015878) Human Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	Antizyme inhibitor 1 (AZIN1) (NM_015878) Human Tagged ORF Clone Lentiviral Particle
Symbol:	Antizyme inhibitor 1
Synonyms:	AZI; AZI1; AZIA1; OAZI; OAZIN; ODC1L
Mammalian Cell Selection:	None
Vector:	pLenti-C-mGFP (PS100071)
Tag:	mGFP
ACCN:	NM_015878
ORF Size:	1344 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC200022).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_015878.4</a>
RefSeq Size:	4385 bp
RefSeq ORF:	1347 bp
Locus ID:	51582
UniProt ID:	<a href="#">O14977</a>
Cytogenetics:	8q22.3
Domains:	Orn_Arg_deC_N
Protein Families:	Druggable Genome



[View online »](#)

**MW:** 49.5 kDa

**Gene Summary:** The protein encoded by this gene belongs to the antizyme inhibitor family, which plays a role in cell growth and proliferation by maintaining polyamine homeostasis within the cell. Antizyme inhibitors are homologs of ornithine decarboxylase (ODC, the key enzyme in polyamine biosynthesis) that have lost the ability to decarboxylase ornithine; however, retain the ability to bind to antizymes. Antizymes negatively regulate intracellular polyamine levels by binding to ODC and targeting it for degradation, as well as by inhibiting polyamine uptake. Antizyme inhibitors function as positive regulators of polyamine levels by sequestering antizymes and neutralizing their effect. This gene encodes antizyme inhibitor 1, the first member of this gene family that is ubiquitously expressed, and is localized in the nucleus and cytoplasm. Overexpression of antizyme inhibitor 1 gene has been associated with increased proliferation, cellular transformation and tumorigenesis. Gene knockout studies showed that homozygous mutant mice lacking functional antizyme inhibitor 1 gene died at birth with abnormal liver morphology. RNA editing of this gene, predominantly in the liver tissue, has been linked to the progression of hepatocellular carcinoma. Alternatively spliced transcript variants have been described for this gene. [provided by RefSeq, Sep 2014]