

## Product datasheet for **RC200064**

### ITGB3BP (NM\_014288) Human Tagged ORF Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** ITGB3BP (NM\_014288) Human Tagged ORF Clone  
**Tag:** Myc-DDK  
**Symbol:** ITGB3BP  
**Synonyms:** CENP-R; CENPR; HSU37139; NRIF3; TAP20  
**Mammalian Cell Selection:** Neomycin  
**Vector:** pCMV6-Entry (PS100001)  
**E. coli Selection:** Kanamycin (25 ug/mL)  
**ORF Nucleotide Sequence:** >RC200064 ORF sequence  
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGATCGCC**

ATGCCTGTAAAAGATCACTGAAGTTGGATGGTCTGTTAGAAGAAAATTCATTTGATCCTTCAAAAATCA  
 CAAGGAAGAAAAGTGTATAACTTATTCTCCAACAAGTGGAACTGTCAAATGAGTCTATTTGCTTCTCC  
 CACAAGTTCTGAAGAGCAAAAGCACAGAAATGGACTATCAAATGAAAAGAGAAAAAATGAATCACCCC  
 AGTTAACTGAAAGCAAAGAATCTACAACAAAAGACAATGATGAATTCATGATGTTGCTATCAAAGTTG  
 AGAAATTGTCAGAAGAAATCATGGAGATAATGCAAAATTAAGTAGTATACAGGCTTTGGAGGGCAGTAG  
 AGAGCTTGAAAATCTCATTGGAATCTCCTGTGCATCACATTTCTTAAAAAGAGAAATGCAGAAAACCAA  
 GAACTAATGACAAAAGTGAATAACAAAACACTGTTTAAAAAGAGTACAGGACTTCTCACAAAGCATCAC  
 GTCATCTTGACAGCTATGAATTCCTTAAAGCCATTTTAAAC

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
 ACAAGGATGACGACGATAAGGTTTAA

**Protein Sequence:** >RC200064 protein sequence  
 Red=Cloning site Green=Tags(s)

MPVKRSLKLDGELLENSFDPSKITRKSVITYSPTTGTCQMSLFASPTSSEEQKHRNGLSNEKRKLNHP  
 SLTESKESTTKDNDFMMLL SKVEKLSEEIMEIMQNLSSIQALEGSRELENLIGISCASHFLKREMOKTK  
 ELMTKVKNKQLFEKSTGLPHKASRHLDSEYFLKAILN

**TRTRPLEQKLI**SEEDLAANDILDYKDDDDKV


**Chromatograms:** [https://cdn.origene.com/chromatograms/mk6052\\_b10.zip](https://cdn.origene.com/chromatograms/mk6052_b10.zip)



[View online »](#)

**Restriction Sites:** SgfI-MluI

**Cloning Scheme:**



**ACCN:** NM\_014288

**ORF Size:** 531 bp

**OTI Disclaimer:** The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [NM\\_014288.5](#)

**RefSeq Size:** 1019 bp

**RefSeq ORF:** 534 bp

**Locus ID:** 23421

**UniProt ID:** [Q13352](#)


**Cytogenetics:** 1p31.3

**Protein Families:** Druggable Genome, Stem cell - Pluripotency, Transcription Factors


**MW:** 20.2 kDa

**Gene Summary:** This gene encodes a transcriptional coregulator that binds to and enhances the activity of members of the nuclear receptor families, thyroid hormone receptors and retinoid X receptors. This protein also acts as a corepressor of NF-kappaB-dependent signaling. This protein induces apoptosis in breast cancer cells through a caspase 2-mediated signaling pathway. This protein is also a component of the centromere-specific histone H3 variant nucleosome associated complex (CENP-NAC) and may be involved in mitotic progression by recruiting the histone H3 variant CENP-A to the centromere. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Sep 2011]


**Product images:**




Circular map for RC200064



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY ITGB3BP (Cat# RC200064, Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ITGB3BP (Cat# [TA803458]). Positive lysates [LY402304] (100ug) and [LC402304] (20ug) can be purchased separately from OriGene.



Western blot validation of overexpression lysate (Cat# [LY402304]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC200064 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).



Coomassie blue staining of purified ITGB3BP protein (Cat# [TP300064]). The protein was produced from HEK293T cells transfected with ITGB3BP cDNA clone (Cat# RC200064) using MegaTran 2.0 (Cat# [TT210002]).