

## Product datasheet for RC200108L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## IDH3B (NM\_006899) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type: Lentiviral Particles

Product Name: IDH3B (NM\_006899) Human Tagged ORF Clone Lentiviral Particle

Symbol: IDH3E Synonyms: RP46

Mammalian Cell Puromycin

Selection:

-

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM\_006899

 ORF Size:
 1155 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC200108).

\_\_\_\_

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 006899.3

RefSeq Size: 1561 bp
RefSeq ORF: 1158 bp
Locus ID: 3420
UniProt ID: O43837
Cytogenetics: 20p13
Domains: isodh

**Protein Pathways:** Citrate cycle (TCA cycle), Metabolic pathways





ORIGENE

**MW:** 42.2 kDa

**Gene Summary:** 

Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These enzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate dehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydrogenases, one of which is mitochondrial and the other predominantly cytosolic. NAD(+)-dependent isocitrate dehydrogenases catalyze the allosterically regulated rate-limiting step of the tricarboxylic acid cycle. Each isozyme is a heterotetramer that is composed of two alpha subunits, one beta subunit, and one gamma subunit. The protein encoded by this gene is the beta subunit of one isozyme of NAD(+)-dependent isocitrate dehydrogenase. Multiple alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Sep 2016]