

## Product datasheet for RC200114L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## C20orf20 (MRGBP) (NM 018270) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

Product Name: C20orf20 (MRGBP) (NM 018270) Human Tagged ORF Clone Lentiviral Particle

Symbol: C20orf20

Synonyms: C20orf20; Eaf7; MRG15BP; URCC4

**Mammalian Cell** 

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM 018270

ORF Size: 612 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC200114).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 018270.3

 RefSeq Size:
 1659 bp

 RefSeq ORF:
 615 bp

 Locus ID:
 55257

 UniProt ID:
 Q9NV56

 Cytogenetics:
 20q13.33

**Protein Families:** Transcription Factors

MW: 22.4 kDa







## **Gene Summary:**

Component of the NuA4 histone acetyltransferase (HAT) complex which is involved in transcriptional activation of select genes principally by acetylation of nucleosomal histones H4 and H2A. This modification may both alter nucleosome - DNA interactions and promote interaction of the modified histones with other proteins which positively regulate transcription. This complex may be required for the activation of transcriptional programs associated with oncogene and proto-oncogene mediated growth induction, tumor suppressor mediated growth arrest and replicative senescence, apoptosis, and DNA repair. NuA4 may also play a direct role in DNA repair when recruited to sites of DNA damage. [UniProtKB/Swiss-Prot Function]