

## Product datasheet for RC200117L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Histone H2A.J (H2AFJ) (NM\_177925) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: Histone H2A. | (H2AF|) (NM 177925) Human Tagged ORF Clone Lentiviral Particle

Symbol: Histone H2A.J

Synonyms: H2AFJ

Mammalian Cell Pu

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 177925

ORF Size: 387 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC200117).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 177925.1

 RefSeq Size:
 3699 bp

 RefSeq ORF:
 390 bp

 Locus ID:
 55766

 UniProt ID:
 Q9BTM1

 Cytogenetics:
 12p12.3

**Protein Pathways:** Systemic lupus erythematosus

**MW:** 14 kDa





## **Gene Summary:**

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is located on chromosome 12 and encodes a replication-independent histone that is a variant H2A histone. The protein is divergent at the C-terminus compared to the consensus H2A histone family member. This gene also encodes an antimicrobial peptide with antibacterial and antifungal activity.[provided by RefSeq, Oct 2015]