

## Product datasheet for RC200160L4

### C9orf95 (NMRK1) (NM\_017881) Human Tagged Lenti ORF Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	C9orf95 (NMRK1) (NM_017881) Human Tagged Lenti ORF Clone
Tag:	mGFP
Symbol:	C9orf95
Synonyms:	bA235O14.2; C9orf95; NRK1
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-mGFP-P2A-Puro (PS100093)
E. coli Selection:	Chloramphenicol (34 ug/mL)
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC200160).
Restriction Sites:	SgfI-MluI
Cloning Scheme:	

Cloning sites used for ORF Shuttling:



\* The last codon before the Stop codon of the ORF.


ACCN:	NM_017881
ORF Size:	597 bp



[View online »](#)

<b>OTI Disclaimer:</b>	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
<b>OTI Annotation:</b>	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_017881.1</a>
<b>RefSeq Size:</b>	1207 bp
<b>RefSeq ORF:</b>	600 bp
<b>Locus ID:</b>	54981
<b>UniProt ID:</b>	<a href="#">Q9NWW6</a>
<b>Cytogenetics:</b>	9q21.13
<b>Protein Pathways:</b>	Nicotinate and nicotinamide metabolism
<b>MW:</b>	23.2 kDa
<b>Gene Summary:</b>	Nicotinamide adenine dinucleotide (NAD <sup>+</sup> ) is essential for life in all organisms, both as a coenzyme for oxidoreductases and as a source of ADP-ribosyl groups used in various reactions. Nicotinic acid and nicotinamide, collectively known as niacin, are the vitamin precursors of NAD <sup>+</sup> . Nicotinamide riboside kinases, such as NRK1, function to synthesize NAD <sup>+</sup> through nicotinamide mononucleotide using nicotinamide riboside as the precursor (Bieganski and Brenner, 2004 [PubMed 15137942]).[supplied by OMIM, Mar 2008]

Product images:



Circular map for RC200160L4