

Product datasheet for RC200204L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NUDT5 (NM_014142) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: NUDT5 (NM 014142) Human Tagged ORF Clone Lentiviral Particle

Symbol: NUDT5

Synonyms: hNUDT5; YSA1; YSA1H; YSAH1

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_014142

ORF Size: 657 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC200204).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 014142.2

RefSeq Size: 1224 bp
RefSeq ORF: 660 bp
Locus ID: 11164
UniProt ID: Q9UKK9
Cytogenetics: 10p14
Domains: NUDIX

Protein Pathways: Purine metabolism





ORIGENE

MW: 24.3 kDa

Gene Summary: This gene belongs to the Nudix (nucleoside diphosphate linked moiety X) hydrolase

superfamily. The encoded enzyme catalyzes the hydrolysis of modified nucleoside diphosphates, including ADP-ribose (ADPR) and 8-oxoGua-containing 8-oxo-dADP and 8-oxo-dGDP. Protein-bound ADP ribose can be hazardous to the cell because it can modify some amino acid residues, resulting in the inhibition of ATP-activated potassium channels. 8-oxoGua is an oxidized form of guanine that can potentially alter genetic information by pairing with adenine and cytosine in RNA. Presence of 8-oxoGua in RNA results in formation of abnormal proteins due to translational errors. [provided by RefSeq, Aug 2013]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US