

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for RC200210L2V

## AKR1C3 (NM\_003739) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

Product Type:	Lentiviral Particles
Product Name:	AKR1C3 (NM_003739) Human Tagged ORF Clone Lentiviral Particle
Symbol:	AKR1C3
Synonyms:	DD3; DDX; HA1753; HAKRB; HAKRe; hluPGFS; HSD17B5; PGFS
Mammalian Cell Selection:	None
Vector:	pLenti-C-mGFP (PS100071)
Tag:	mGFP
ACCN:	NM_003739
ORF Size:	969 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC200210).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<u>NM 003739.4</u>
RefSeq Size:	1251 bp
RefSeq ORF:	972 bp
Locus ID:	8644
UniProt ID:	<u>P42330</u>
Cytogenetics:	10p15.1
Protein Families:	Druggable Genome
Protein Pathways:	Arachidonic acid metabolism, Metabolism of xenobiotics by cytochrome P450



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	AKR1C3 (NM_003739) Human Tagged ORF Clone Lentiviral Particle – RC200210L2V
MW:	36.9 kDa
Gene Summary:	This gene encodes a member of the aldo/keto reductase superfamily, which consists of more than 40 known enzymes and proteins. These enzymes catalyze the conversion of aldehydes and ketones to their corresponding alcohols by utilizing NADH and/or NADPH as cofactors. The enzymes display overlapping but distinct substrate specificity. This enzyme catalyzes the reduction of prostaglandin (PG) D2, PGH2 and phenanthrenequinone (PQ), and the oxidation of 9alpha,11beta-PGF2 to PGD2. It may play an important role in the pathogenesis of allergic diseases such as asthma, and may also have a role in controlling cell growth and/or differentiation. This gene shares high sequence identity with three other gene members and is clustered with those three genes at chromosome 10p15-p14. Three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Dec 2011]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US