

## Product datasheet for RC200320L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## APEX2 (NM\_014481) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** APEX2 (NM\_014481) Human Tagged ORF Clone Lentiviral Particle

Symbol: APEX2

**Synonyms:** APE2; APEXL2; XTH2; ZGRF2

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_014481 **ORF Size:** 1554 bp

**ORF Nucleotide** 

- - - -

Sequence:
OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC200320).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 014481.2

 RefSeq Size:
 2095 bp

 RefSeq ORF:
 1557 bp

 Locus ID:
 27301

 UniProt ID:
 Q9UBZ4

 Cytogenetics:
 Xp11.21

Protein Families: Druggable Genome
Protein Pathways: Base excision repair




MW:

57.4 kDa

**Gene Summary:** 

Apurinic/apyrimidinic (AP) sites occur frequently in DNA molecules by spontaneous hydrolysis, by DNA damaging agents or by DNA glycosylases that remove specific abnormal bases. AP sites are pre-mutagenic lesions that can prevent normal DNA replication so the cell contains systems to identify and repair such sites. Class II AP endonucleases cleave the phosphodiester backbone 5' to the AP site. This gene encodes a protein shown to have a weak class II AP endonuclease activity. Most of the encoded protein is located in the nucleus but some is also present in mitochondria. This protein may play an important role in both nuclear and mitochondrial base excision repair. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Nov 2012]

## **Product images:**



[RC200320L4] was used to prepare Lentiviral particles using [TR30037] packaging kit. HEK293T cells were transduced with RC200320L4V particle to overexpress human APEX2-mGFP fusion protein.