

Product datasheet for **RC200353**

PSMB8 (NM_004159) Human Tagged ORF Clone

Product data:

Product Type: Expression Plasmids
Product Name: PSMB8 (NM_004159) Human Tagged ORF Clone
Tag: Myc-DDK
Symbol: PSMB8
Synonyms: ALDD; D6S216; D6S216E; JMP; LMP7; NKJO; PRAAS1; PSMB5i; RING10
Mammalian Cell Selection: Neomycin
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
ORF Nucleotide Sequence: >RC200353 representing NM_004159
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGCTCATAGGAACCCCAACCCCGCTGACACTACTCCAGCTCCTGGCTGACTTCTAGTCTTCTGGTTG
AAGCTGCGCCTTTAGATGACACGACCCTACCCACCCTGTTTCCAGCGGATGCCCGGCCCTGGAGCCAC
AGAATTCTCCAGTCCCTGGGTGGGACGGAGAAAGAACGTTTCAGATTGAGATGGCCCATGGCACCACC
ACGCTCGCCTTCAAGTTCAGCATGGAGTATTGCAGCAGTGGATTCTCGGGCCTCAGCTGGTCTTACA
TTAGTGCCTTACGGGTGAACAAGGTGATTGAGATTAACCCTTACCTGCTTGGCACCATGTCTGGCTGTGC
AGCAGACTGTGACTACTGGGAGCGCTGCTGGCCAAGGAATGCAGGCTGTACTATCTGCGAAATGGAGAA
CGTATTTTCAGTGTCCGACGCTCAAGCTGCTGTCCAACATGATGTGCCAGTACCGGGGCATGGGCCTCT
CTATGGGCAGTATGATCTGTGGCTGGGATAAGAAGGGTCTGGACTCTACTACGTGGATGAACATGGGAC
TCGGCTCTCAGGAAATATGTTCTCCACGGGTAGTGGGAACACTTATGCCTACGGGGTCAATGGACAGTGGC
TATCGGCCTAATCTTAGCCCTGAAGAGGCTATGACCTTGGCCGAGGGCTATTGCTTATGCCACTACA
GAGACAGCTATTCTGGAGCGTTGTCAATATGTACCACATGAAGGAAGATGGTTGGGTGAAAGTAGAAAG
TACAGATGTCAGTGACCTGCTGCACCAGTACCGGAAGCCAATCAA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA



Protein Sequence: >RC200353 representing NM_004159
 Red=Cloning site Green=Tags(s)

MLIGTPTPRDTPSSWL TSSLLVEAAPLDDTTLTPVSSGCPGLEPTEFFQSLGGDGERNVQIEMAHGTT
 TLAFKFQHGVIAAVDSRASAGSYISALRVNKVIEINPYLLGTMSGCAADCQYWERLLAKECRLYYLRNGE
 RISVSAASKLLSNMMCQYRGMGLSMGSMICGWDKKGPGLYYVDEHGTRLSGNMFSTGSGNTYAYGVMDSG
 YRPNLSPEEAYDLGRRAIAYATHRDSYSGGVNMYHMKEDGWVKVESTDVSDDLHQYREANQ


TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mg2875_g08.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:

Cloning sites used for ORF Shuttling:



* The last codon before the Stop codon of the ORF

ACCN: NM_004159

ORF Size: 816 bp


OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.


Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	NM_004159.5
RefSeq Size:	1602 bp
RefSeq ORF:	819 bp
Locus ID:	5696
UniProt ID:	P28062
Cytogenetics:	6p21.32
Protein Families:	Druggable Genome, Protease
Protein Pathways:	Proteasome
MW:	29.6 kDa
Gene Summary:	<p>The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the proteasome B-type family, also known as the T1B family, that is a 20S core beta subunit. This gene is located in the class II region of the MHC (major histocompatibility complex). Expression of this gene is induced by gamma interferon and this gene product replaces catalytic subunit 3 (proteasome beta 5 subunit) in the immunoproteasome. Proteolytic processing is required to generate a mature subunit. Two alternative transcripts encoding two isoforms have been identified; both isoforms are processed to yield the same mature subunit. [provided by RefSeq, Jul 2008]</p>


Product images:



Circular map for RC200353



Western blot validation of overexpression lysate (Cat# [LY418176]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC200353 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).



Coomassie blue staining of purified PSMB8 protein (Cat# [TP300353]). The protein was produced from HEK293T cells transfected with PSMB8 cDNA clone (Cat# RC200353) using MegaTran 2.0 (Cat# [TT210002]).