

## Product datasheet for RC200411L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## TGFBI (NM\_000358) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: TGFBI (NM 000358) Human Tagged ORF Clone Lentiviral Particle

Symbol: TGFB

Synonyms: BIGH3; CDB1; CDG2; CDGG1; CSD1; CSD2; CSD3; EBMD; LCD1

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM\_000358

ORF Size: 2049 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC200411).

Sequence:
OTI Disclaimer:

**Domains:** 

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 000358.1

 RefSeq Size:
 2805 bp

 RefSeq ORF:
 2052 bp

 Locus ID:
 7045

 UniProt ID:
 Q15582

 Cytogenetics:
 5q31.1

**Protein Families:** Druggable Genome, Secreted Protein, Transmembrane

FAS1





## TGFBI (NM\_000358) Human Tagged ORF Clone Lentiviral Particle - RC200411L1V

**MW:** 74.7 kDa

**Gene Summary:** 

This gene encodes an RGD-containing protein that binds to type I, II and IV collagens. The RGD motif is found in many extracellular matrix proteins modulating cell adhesion and serves as a ligand recognition sequence for several integrins. This protein plays a role in cell-collagen interactions and may be involved in endochondrial bone formation in cartilage. The protein is induced by transforming growth factor-beta and acts to inhibit cell adhesion. Mutations in this gene are associated with multiple types of corneal dystrophy. [provided by RefSeq, Jul 2008]