

## Product datasheet for RC200420L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## CX3CL1 (NM\_002996) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** CX3CL1 (NM\_002996) Human Tagged ORF Clone Lentiviral Particle

Symbol: CX3CL1

Synonyms: ABCD-3; C3Xkine; CXC3; CXC3C; fractalkine; neurotactin; NTN; NTT; SCYD1

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-mGFP (PS100071)

Tag: mGFP

**ACCN:** NM\_002996 **ORF Size:** 1191 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC200420).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 002996.3

 RefSeq Size:
 3304 bp

 RefSeq ORF:
 1194 bp

 Locus ID:
 6376

 UniProt ID:
 P78423

 Cytogenetics:
 16q21

Domains: IL8

**Protein Families:** Druggable Genome, Secreted Protein, Transmembrane





## CX3CL1 (NM\_002996) Human Tagged ORF Clone Lentiviral Particle - RC200420L2V

**Protein Pathways:** Chemokine signaling pathway, Cytokine-cytokine receptor interaction

MW: 42.7 kDa

**Gene Summary:** This gene belongs to the CX3C subgroup of chemokines, characterized by the number of

amino acids located between the conserved cysteine residues. This is the only member of the CX3C subgroup, which contains three amino acids between cysteine residues, resulting in a Cys-X-X-Cys configuration. The encoded protein contains an extended mucin-like stalk with a chemokine domain on top, and exists in both a membrane-anchored form where it acts as a binding molecule, or, in soluble form, as a chemotactic cytokine. The mature form of this protein can be cleaved at the cell surface, yielding different soluble forms that can interact with the G-protein coupled receptor, C-X3-C motif chemokine receptor 1 gene product. This gene plays a role in a wide range of diseases, including cancer, vasculitis, neuropathies, atherosclerosis, inflammatory diseases, and in human immunodeficiency virus infections.

[provided by RefSeq, Sep 2017]