

## Product datasheet for RC200423L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## RPL38 (NM\_001035258) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

Product Name: RPL38 (NM 001035258) Human Tagged ORF Clone Lentiviral Particle

Symbol: RPL38
Synonyms: L38

Mammalian Cell Pur

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

**ACCN:** NM\_001035258

ORF Size: 210 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC200423).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 001035258.1

 RefSeq Size:
 376 bp

 RefSeq ORF:
 213 bp

 Locus ID:
 6169

 UniProt ID:
 P63173

 Cytogenetics:
 17q25.1

**Protein Families:** Druggable Genome

**Protein Pathways:** Ribosome





ORIGENE

**MW:** 8.2 kDa

**Gene Summary:** 

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L38E family of ribosomal proteins. It is located in the cytoplasm. Alternative splice variants have been identified, both encoding the same protein. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome, including one located in the promoter region of the type 1 angiotensin II receptor gene. [provided by RefSeq, Jul 2008]