

Product datasheet for RC200426L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RFC4 (NM_002916) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: RFC4 (NM_002916) Human Tagged ORF Clone Lentiviral Particle

Symbol: RFC4

Synonyms: A1; RFC37

Mammalian Cell Puromycin

Selection:

Vector:

pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 002916

ORF Size: 1089 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC200426).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 002916.3

 RefSeq Size:
 1427 bp

 RefSeq ORF:
 1092 bp

 Locus ID:
 5984

 UniProt ID:
 P35249

 Cytogenetics:
 3q27.3

Domains: AAA, AAA

Protein Families: Druggable Genome, Stem cell - Pluripotency





RFC4 (NM_002916) Human Tagged ORF Clone Lentiviral Particle - RC200426L3V

Protein Pathways: DNA replication, Mismatch repair, Nucleotide excision repair

MW: 39.7 kDa

Gene Summary: The elongation of primed DNA templates by DNA polymerase delta and DNA polymerase

epsilon requires the accessory proteins proliferating cell nuclear antigen (PCNA) and replication factor C (RFC). RFC, also named activator 1, is a protein complex consisting of five distinct subunits of 140, 40, 38, 37, and 36 kD. This gene encodes the 37 kD subunit. This subunit forms a core complex with the 36 and 40 kDa subunits. The core complex possesses DNA-dependent ATPase activity, which was found to be stimulated by PCNA in an in vitro system. Alternatively spliced transcript variants encoding the same protein have been

reported. [provided by RefSeq, Jul 2008]