

## Product datasheet for RC200444L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PDZK1 (NM\_002614) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

Product Name: PDZK1 (NM\_002614) Human Tagged ORF Clone Lentiviral Particle

Symbol: PDZK1

Synonyms: CAP70; CLAMP; NHERF-3; NHERF3; PDZD1

**Mammalian Cell** 

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM\_002614

 ORF Size:
 1557 bp

**ORF Nucleotide** 

Sequence:

The ORF insert of this clone is exactly the same as(RC200444).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 002614.3</u>

 RefSeq Size:
 2301 bp

 RefSeq ORF:
 1560 bp

 Locus ID:
 5174

 UniProt ID:
 Q5T2W1

 Cytogenetics:
 1q21.1

Cytogenetics: 1q21

Domains: PDZ

**MW:** 57.1 kDa







## **Gene Summary:**

This gene encodes a PDZ domain-containing scaffolding protein. PDZ domain-containing molecules bind to and mediate the subcellular localization of target proteins. The encoded protein mediates the localization of cell surface proteins and plays a critical role in cholesterol metabolism by regulating the HDL receptor, scavenger receptor class B type 1. Single nucleotide polymorphisms in this gene may be associated with metabolic syndrome, and overexpression of this gene may play a role in drug resistance of multiple myeloma. Pseudogenes of this gene are located on the long arm of chromosome 1. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Jan 2011]