

Product datasheet for RC200446L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OGDH (NM_002541) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: OGDH (NM_002541) Human Tagged ORF Clone Lentiviral Particle

Symbol: OGDH

Synonyms: AKGDH; E1k; KGD1; OGDC; OGDH2

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM_002541

 ORF Size:
 3069 bp

ORF Nucleotide

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(RC200446).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 002541.2

 RefSeq Size:
 4319 bp

 RefSeq ORF:
 3072 bp

 Locus ID:
 4967

 UniProt ID:
 Q02218

Cytogenetics: 7p13

Domains: E1_dehydrog, transket_pyr

Protein Families: Druggable Genome





OGDH (NM_002541) Human Tagged ORF Clone Lentiviral Particle - RC200446L1V

Protein Pathways: Citrate cycle (TCA cycle), Lysine degradation, Metabolic pathways, Tryptophan metabolism

MW: 115.9 kDa

Gene Summary: This gene encodes one subunit of the 2-oxoglutarate dehydrogenase complex. This complex

catalyzes the overall conversion of 2-oxoglutarate (alpha-ketoglutarate) to succinyl-CoA and CO(2) during the Krebs cycle. The protein is located in the mitochondrial matrix and uses

thiamine pyrophosphate as a cofactor. A congenital deficiency in 2-oxoglutarate dehydrogenase activity is believed to lead to hypotonia, metabolic acidosis, and

hyperlactatemia. Alternative splicing results in multiple transcript variants encoding distinct

isoforms.[provided by RefSeq, Sep 2009]