

## Product datasheet for RC200454L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## MATK (NM\_139354) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** MATK (NM\_139354) Human Tagged ORF Clone Lentiviral Particle

Symbol: MATK

Synonyms: CHK; CTK; HHYLTK; HYL; HYLTK; Lsk

**Mammalian Cell** 

viairiirianari een

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_139354 **ORF Size:** 1521 bp

**ORF Nucleotide** 

- -

Sequence:

The ORF insert of this clone is exactly the same as(RC200454).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 139354.2

 RefSeq Size:
 1940 bp

 RefSeq ORF:
 1401 bp

 Locus ID:
 4145

 UniProt ID:
 P42679

 Cytogenetics:
 19p13.3

**Protein Families:** Druggable Genome, Protein Kinase, Stem cell - Pluripotency

**MW:** 56.5 kDa







## **Gene Summary:**

The protein encoded by this gene has amino acid sequence similarity to Csk tyrosine kinase and has the structural features of the CSK subfamily: SRC homology SH2 and SH3 domains, a catalytic domain, a unique N terminus, lack of myristylation signals, lack of a negative regulatory phosphorylation site, and lack of an autophosphorylation site. This protein is thought to play a significant role in the signal transduction of hematopoietic cells. It is able to phosphorylate and inactivate Src family kinases, and may play an inhibitory role in the control of T-cell proliferation. This protein might be involved in signaling in some cases of breast cancer. Three alternatively spliced transcript variants that encode different isoforms have been described for this gene. [provided by RefSeq, Jul 2008]